

ABSTRACT SUBMISSION COMPLETE

Abstract Information > Abstract Review > **Abstract Submission**

Confirmation #: 27028

Thank you for submitting an abstract to the ASLO Meeting 2015 in Granada, Spain. This is confirmation that your submission has been processed and is complete. Please print this page as a record of your transaction.

You will receive a confirmation message to the registration e-mail address shortly. Please contact the ASLO Meeting 2015 in Granada,

Spain Management Office by telephone (+1 254-776-3550) or e-mail (aslogranada2015@sgmeet.com) if you do not receive a confirmation message within 24 hours.

We encourage you to visit the conference web site often for updates and the most current information as it becomes available. We look forward to seeing you!

If your abstract submission completes your registration, please click below to return to the conference home page:

[Return to Meeting Home Page](#)

2015 ASLO Aquatic Sciences Meeting Abstract
Submission Confirmation

Thank you for submitting your abstract for the ASLO Meeting 2015 in Granada, Spain. We look forward to seeing you!

We encourage you to visit the conference website often for updates and the most current information as it becomes available - <http://www.sgmeet.com/aslo/granada2015/>.

Date: 10/10/2014 10:01:32 AM

Abstract ID: 27028

To review/edit your abstract, please go to - <https://www.sgmeet.com/aslo/granada2015/userlogin.asp>

LoginID: mluz.fernandez@ba.ieo.es

Password: 16798

Your abstract details are as follows:

Title: VERTICAL BIOGEOGRAPHICAL OVERVIEW OF THE ZOOPLANKTON COMMUNITY
ACROSS THE ATLANTIC, PACIFIC AND INDIAN OCEAN (35°N-40°S)

Text: From December 2010 to June 2011 the mesozooplankton has been sampled at the deep Atlantic, Indian and Pacific Ocean (*). From surface to 3000 m depth a HYDROBIOSS multinet was used distinguishing the epipelagic (77%), mesopelagic (11%) and bathypelagic zones (11%), where five layers were usually sampled (0-200, 200-500, 500-1000, 1000-2000 and 2000-3000 m depth). Among the three oceans, no large differences on abundance were found when taxonomic groups were considered, being always depth the most important factor affecting the vertical zooplankton distribution. The zooplankton abundance strongly decreased with depth and very low abundance was found at deeper waters. Very irregular spatial distribution was observed all across the three oceans, finding the lowest abundance in the south and western Pacific region. Copepods were always the most abundant contributors of the zooplankton community (84%) and more than 260 species identified. They were followed by chaetognaths (5%), siphonophores (3%), ostracods (2%) and euphausiids (1%). In a biogeographical overview, the vertical distribution of the most abundant copepods is analyzed, finding the largest copepods at deeper stratum, where small cosmopolitan copepods were also found.

Presentation Choice: Oral

Session Choice 1: 008

Session Choice 2: 017

Confirmation #: 27028

For comments or questions, please contact:

2015 ASLO Aquatic Sciences Meeting Management 5400 Bosque Blvd., Suite
680 Waco, Texas 76710-4446 254-776-3550 Phone

254-776-3767 Fax

aslogranada2015@sgmeet.com

<http://www.sgmeet.com/aslo/granada2015/>

Important: Do not respond directly to this message. This message is for confirmation only. For comments or questions, please contact the 2015 ASLO Aquatic Sciences Meeting Management Office.